

# Universally Measurable Sets I

Paul Larson

Department of Mathematics  
Miami University  
Oxford, Ohio 45056  
larsonpb@miamioh.edu

February 2, 2026

## Measures

Universally  
measurable setsUniversally  
null sets

Category

Reclaw's  
TheoremCichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

Given a set  $X$ , a  $\sigma$ -algebra on  $X$  is a nonempty collection of subsets of  $X$  which is closed under countable unions and complements.

A *measure* on  $X$  is a function  $\mu$  from some  $\sigma$ -algebra on  $X$  to  $[0, \infty]$ , such that  $\mu(\emptyset) = 0$  and, whenever  $A_n$  ( $n \in \omega$ ) are disjoint members of the domain of  $\mu$ ,

$$\mu\left(\bigcup_{n \in \omega} A_n\right) = \sum_{n \in \omega} \mu(A_n).$$

## Measures

Universally  
measurable sets

Universally  
null sets

Category

Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

A measure  $\mu$  on a set  $X$  is

- *finite* if  $\mu(X)$  is finite;
- $\sigma$ -*finite* if  $X$  is a countable union of sets with finite measure;
- a *probability measure* if  $\mu(X) = 1$ ;
- *atomless* if  $\mu(a) = 0$  for all finite  $a \subseteq X$ .

## Measures

Universally  
measurable sets

Universally  
null sets

Category

Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

Given a topological space  $(X, \tau)$ , the Borel subsets of  $X$  are the members of the smallest  $\sigma$ -algebra on  $X$  containing the open sets.

We will write  $\text{Borel}(X)$  for the set of Borel subsets of  $X$ .

A *Borel measure* on  $(X, \tau)$  is a measure whose domain contains the Borel sets.

## Measures

Universally  
measurable sets

Universally  
null sets

Category

Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

A measure  $\mu$  on a set  $X$  is *complete* if  $A \in \text{dom}(\mu)$  whenever there exist  $B, N \in \text{dom}(\mu)$  such that  $A \triangle B \subseteq N$  and  $\mu(N) = 0$  (in which case  $\mu(A) = \mu(B)$ ).

The *completion* of a measure is the smallest complete measure containing it (i.e., the intersection of all the complete measures containing it).

## Measures

Universally  
measurable sets

Universally  
null sets

Category

Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

- Lebesgue measure on  $\mathbb{R}$  ( $\sigma$ -finite and atomless; its restriction to  $[0, 1]$  is a probability measure;  $(a, b)$  gets measure  $b - a$ );
- Haar measure on  $2^\omega$  (an atomless probability measure;  $[s]$  gets measure  $2^{-|s|}$ ).

## Measures

Universally  
measurable setsUniversally  
null sets

Category

Reclaw's  
TheoremCichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

A function between topological spaces is *Borel* if the preimage of each open set is Borel.

If  $\mu$  is a Borel measure on a topological space  $X$ , and

$$f: X \rightarrow \mathbb{R}^{\geq 0}$$

is a Borel function, then

$$\nu(A) = \int_A f(x) d\mu$$

defines another Borel measure on  $X$ .

## Measures

Universally  
measurable sets

Universally  
null sets

Category

Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

**Definition.** A subset of  $\mathbb{R}$  (or, more generally, of any Polish space) is *analytic* (or  $\Sigma_1^1$ ) if it is a continuous image of a Borel set, and *coanalytic* (or  $\Pi_1^1$ ) if it is the complement of an analytic set.

By a classical theorem of Lusin, analytic subsets of  $\mathbb{R}$  or  $2^\omega$  are Lebesgue/Haar measurable (so coanalytic sets are too).

Four classical ways in which nonmeasurable sets arise (using the Axiom of Choice):

- Choosing a member from each equivalence class of  $\mathbb{R}/\mathbb{Q}$  (forming a Vitali set).
- Fubini's Theorem, which can be used to show that no wellordering of  $\mathbb{R}$  is Lebesgue measurable.
- Nonprincipal ultrafilters on  $\omega$ .
- Rapid filters.

**Definition.** A subset of a topological space is *universally measurable* if it is measurable with respect to (i.e., in the domain of) every complete  $\sigma$ -finite Borel measure on the space.

We will write  $\text{UMeas}(X)$  for the set of universally measurable subsets of a topological space  $(X, \tau)$ .

For any topological space  $(X, \tau)$ ,  $\text{UMeas}(X)$  is a  $\sigma$ -algebra containing the Borel sets.

The set of universally measurable subsets of a topological space would not change if one replaced “ $\sigma$ -finite” with “finite” or “probability”.

It would also not change if one added the requirement that the measures be atomless (since a  $\sigma$ -finite measure can have only countably many atoms).\*

A topological space  $(X, \tau)$  is *completely metrizable* if there exists a metric  $d$  on  $X$  inducing  $\tau$  relative to which  $X$  is complete (i.e., such that every  $d$ -Cauchy sequence converges).

A *Polish space* is a completely metrizable separable topological space.

The class of Polish spaces is closed under countable products and taking  $G_\delta$  subsets.

## Examples of Polish spaces:

- $\mathbb{Z}$
- $\mathbb{R}^n$
- $[0, 1]$
- $2^\omega$
- $(2^\omega)^\omega$
- $\omega^\omega$

The *Suslin operation* applied to sets  $A_\sigma$  ( $\sigma \in \omega^{<\omega}$ ) results in the set

$$\bigcup_{x \in \omega^\omega} \bigcap_{n \in \omega} A_{x \upharpoonright n}.$$

The *C-sets* are the members of the smallest  $\sigma$ -algebra containing the open sets and closed under the Suslin operation.

If  $(X, \tau)$  is Polish then  $\text{UMeas}(X)$  is closed under the Suslin operation and therefore contains the *C-sets*.

The questions we consider in these lectures generally have the following form: what can the set of universally measurable subsets of Polish spaces be like in models of ZFC?

# Universally measurable homomorphisms

P.B. Larson

Measures

Universally measurable sets

Universally null sets

Category

Reclaw's Theorem

Cichoń's diagram

Questions

Nonmeager universally null sets

References

Given topological spaces  $X$  and  $Y$ , a function  $f: X \rightarrow Y$  is *universally measurable* if  $f^{-1}[A]$  is universally measurable for every Borel  $A \subseteq Y$ .

**Theorem** (Rosendal) If  $\pi$  is a universally measurable homomorphism from a Polish group to a separable group then  $\pi$  is continuous.

**Theorem** (Christensen) If  $G = \bigcup_{i \in \mathbb{N}} A_i$  is a covering of a Polish group by universally measurable sets, then for every neighborhood  $U$  of the identity there exist  $k \in \mathbb{N}$  and  $g_1, \dots, g_n \in U$  such that

$$\bigcup_{i=1}^n g_i A_k A_k^{-1} g_i^{-1}$$

is a neighborhood of the identity.

# The Isomorphism Theorem (for measures)

Measures

Universally  
measurable setsUniversally  
null sets

Category

Reclaw's  
TheoremCichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

**Theorem.** If  $X$  and  $Y$  are Polish spaces and  $\mu$  and  $\nu$  are atomless Borel probability measures on  $X$  and  $Y$  respectively, then there is a Borel bijection

$$f: X \rightarrow Y$$

such that

$$\mu(I) = \nu(f[I])$$

for all Borel  $I \subseteq X$ .

The Isomorphism Theorem gives us the following:

- For the sorts of questions that we will be asking, it doesn't matter which Polish space we consider (often we use  $2^\omega$ ).
- Since analytic subsets of  $[0, 1]$  are Lebesgue measurable, and a Borel preimage of an analytic set is analytic, analytic subsets of any Polish space are universally measurable.

If

- $\mu$  is a Borel measure on a topological space  $(X, \tau)$ ,
- $(Y, \sigma)$  is a topological space,
- $f: X \rightarrow Y$  is a Borel function,

then  $\mu_f(A) = \mu(f^{-1}[A])$  defines a Borel measure on  $(Y, \sigma)$ .

This is sometimes called the *pushforward* measure induced by  $f$ .

It follows that

- a subset  $A$  of Polish space  $X$  is universally measurable if and only if  $f^{-1}[A]$  is Lebesgue measurable whenever  $f: 2^\omega \rightarrow X$  is a Borel function.
- If  $(X, \tau)$  and  $(Y, \sigma)$  are Polish spaces,  $A \subseteq Y$  is universally measurable and  $f: X \rightarrow Y$  is Borel, then  $f^{-1}[A]$  is universally measurable.
- In a model of ZF in which every set of reals is Lebesgue measurable (such as a Solovay model), every set of reals is universally measurable.

# Universally Null Sets (I)

P.B. Larson

Measures

Universally  
measurable sets

Universally  
null sets

Category

Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

A subset of a topological space is *universally null* if it has measure 0 with respect to every atomless  $\sigma$ -finite Borel measure on the space.

$\text{UNull}(X)$  denotes the set of universally null subsets of  $(X, \tau)$ .

Again, the collection of universally null sets does not change if we require the measure to be finite, or a probability measure.

# Universally Null Sets (II)

P.B. Larson

Measures

Universally  
measurable sets

Universally  
null sets

Category

Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

$\text{UNull}(X)$  is a  $\sigma$ -ideal on  $X$  (i.e., it is closed under subsets and countable unions) containing all countable sets.

$$\text{UNull}(X) \subseteq \text{UMeas}(X)$$

A subset  $A$  of Polish space  $X$  is universally null if and only if  $f^{-1}[A]$  is Lebesgue null whenever

$$f: 2^\omega \rightarrow X$$

is a Borel function such that the preimage of each point is Lebesgue null.

A subset of a topological space is said to be *perfect* if it is closed and nonempty and contains no isolated points.

We will write  $\text{Perf}(X)$  for the set of perfect subsets of  $(X, \tau)$ .

When  $(X, \tau)$  is an uncountable Polish space, each element of  $\text{Perf}(X)$  has cardinality  $2^{\aleph_0}$ .

For each  $P \in \text{Perf}(X)$ , there is a Borel probability measure  $\mu$  on  $X$  with  $\mu(P) = 1$ .

It follows that a universally null subset of a Polish space cannot contain a perfect set.

Every uncountable analytic subset of a Polish space contains a perfect set (so the analytic universally null sets are exactly the countable sets).

**Theorem** (Grzegorek, Ryll-Nardzewski) The (coanalytic) set of wellorderings of  $\omega$  is universally measurable but does not have universally null symmetric difference with any Borel set.

A subset of a topological space is

- *nowhere dense* if its closure does not contain any nonempty open set, and
- *meager* if it is a countable union of nowhere dense sets.

A set has the Baire Property if its symmetric difference with some open set is meager.

By analogy with the measure case, we make the following definitions.

A subset  $A$  of a Polish space  $(X, \tau)$  is *universally categorical* if  $f^{-1}[A]$  has the Baire property whenever  $(Y, \sigma)$  is a Polish space and  $f: Y \rightarrow X$  is Borel.

A subset of a Polish space is *universally meager* if  $f^{-1}[A]$  is meager whenever  $(Y, \sigma)$  is a Polish space and  $f: Y \rightarrow X$  is Borel such that the preimage of each point is meager.

We write  $\text{UCat}(X)$  for the set of universally categorical subsets of  $X$ , and  $\text{UMeag}(X)$  for the set of universally meager sets.

When  $(X, \tau)$  is a Polish space,  $\text{UCat}(X)$  is a  $\sigma$ -algebra containing the analytic sets and closed under the Suslin operation, and  $\text{UMeag}(X)$  is a  $\sigma$ -ideal containing the countable sets.

**Theorem.**(Reclaw) Let  $(X, \tau)$  be a Polish space. A subset of  $X$  which is wellordered by a universally measurable relation on  $X$  is universally null. A subset of  $X$  which is wellordered by a universally categorical relation on  $X$  is universally meager.

The proof uses Fubini's theorem and (its category-analogue) the Kuratowski-Ulam theorem.

# Uncountable universally null and meager sets (I)

P.B. Larson

Measures

Universally  
measurable sets

Universally  
null sets

Category

Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

The relation of mod-finite domination is Borel (so universally measurable and universally categorical) and contains chains of ordertype  $\mathfrak{b}$ .

It follows (from Reclaw's theorem) that

$$\text{UNull}(\omega^\omega) \cap \text{UMeag}(\omega^\omega)$$

contains sets of cardinality  $\mathfrak{b}$ , and (since both properties are closed under taking subsets) at least  $2^{\mathfrak{b}}$  many such sets.

# Uncountable universally null and meager sets (II)

P.B. Larson

Measures

Universally  
measurable sets

Universally  
null sets

Category

Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

Similarly, one show that for any ideal  $\mathcal{I}$  on any Polish space  $P$ , if  $\mathcal{I}$  is generated by sets of some fixed Borel complexity, then there is a set of cardinality

$$\min\{\text{non}(\mathcal{I}), \text{cov}(\mathcal{I})\}$$

which is wellordered by a Borel relation (and is therefore both universally null and universally meager).

# Cichoń's diagram

P.B. Larson

Measures

Universally  
measurable sets

Universally  
null sets

Category


Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References



For any Polish space  $(X, \tau)$ ,  $\text{UMeas}(X)$  and  $\text{UNull}(X)$  are both closed under unions of cardinality less than  $\text{Add}(\mathcal{N})$ .

Similarly,  $\text{UCat}(X)$  and  $\text{UMeag}(X)$  are both closed under unions of cardinality less than  $\text{Add}(\mathcal{M})$ .

**Theorem.** (Grzegorek) There exist a universally null set of cardinality  $\text{non}(\mathcal{N})$  and a universally meager set of cardinality  $\text{non}(\mathcal{M})$ .

One can use this theorem to show that the classes of universally measurable and universally categorical (check) sets are not closed under projections.

Is it consistent with ZFC that every universally measurable set has the property of Baire (in which case it would be universally categorical)?

Is it consistent with ZFC that every universally categorical set is universally measurable?

Is it consistent with ZFC that  $\text{UMeas}(X)$  is the same as the  $C$ -sets? (no way)

Is it consistent with ZFC that every universally measurable set has universally null symmetric difference with some  $C$ -set?

Is it consistent with ZFC that the universally measurable sets are the members of the smallest  $\sigma$ -algebra closed under the Suslin operation and containing the open sets and the universally null sets?

Is it consistent with ZFC that  $\text{UNull}(X) = \text{UMeag}(X)$  (for some/any Polish space  $(X, \tau)$ )?

Recall that the real line is the union of a meager set and a null set, so there are nonmeager null sets and vice versa.

# Nonmeager universally null sets

P.B. Larson

Measures

Universally  
measurable sets

Universally  
null sets

Category

Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

Since universally null sets don't contain perfect sets, a universally null set which has the Baire property must be meager.

**Theorem**(Brendle-Larson) If  $\min\{\mathfrak{d}, \text{non}(\mathcal{N})\} = \text{cof}(\mathcal{M})$ , then there is a nonmeager universally null set of cardinality  $\text{cof}(\mathcal{M})$ .

We build a nonmeager universally null set in the Polish space  $\omega^\omega$ . A key point is that if  $\mu$  is any Borel probability measure on  $\omega^\omega$ , then there is a  $g \in \omega^\omega$  such that  $\mu(B_g) = 1$ , where

$$B_g = \{f : f \leq^* g\}.$$

A tree  $T \subseteq \omega^{<\omega}$  is *superperfect* if, for densely many  $s \in T$ ,

$$\{n : s \frown \langle n \rangle \in T\}$$

is infinite.

We write  $[T]$  for the set of infinite paths through  $T$ .

If  $A \subseteq \omega^\omega$  is meager, then there is a superperfect tree

$$T \subseteq \omega^{<\omega}$$

such that  $A \cap [T] = \emptyset$ .

# Proof of the theorem (I)

P.B. Larson

Measures

Universally  
measurable sets

Universally  
null sets

Category

Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

Let  $\kappa$  be  $\text{cof}(\mathcal{M})$ , which is equal to  $\mathfrak{d}$  by the hypothesis.

Let  $\{A_\alpha : \alpha < \kappa\}$  witness  $\text{cof}(\mathcal{M}) = \kappa$  and let  $\{f_\alpha : \alpha < \kappa\}$  be a dominating family in  $\omega^\omega$ .

For each  $\alpha < \kappa$ , let  $T_\alpha \subseteq \omega^{<\omega}$  be a superperfect tree such that

$$A_\alpha \cap [T_\alpha] = \emptyset.$$

**Claim.** Since  $\mathfrak{d} = \kappa$ , we can pick for each  $\alpha < \kappa$  an  $x_\alpha \in [T_\alpha]$  such that, for all  $\beta < \alpha$ ,  $f_\beta \not\geq^* x_\alpha$ .

# Proof of the theorem (II)

P.B. Larson

Measures

Universally  
measurable sets

Universally  
null sets

Category

Reclaw's  
Theorem

Cichoń's  
diagram

Questions

Nonmeager  
universally null  
sets

References

Since  $X = \{x_\alpha : \alpha < \kappa\}$  is not contained in any  $A_\alpha$ , it is nonmeager.

To see that  $X$  is universally null, let  $\mu$  be a Borel probability measure on  $\omega^\omega$ . Fix  $g \in \omega^\omega$  such that  $\mu(B_g) = 1$ .

Let  $\alpha$  be such that  $g \leq^* f_\alpha$ . Then

$$\{x_\beta : \beta > \alpha\} \cap B_g = \emptyset.$$

Thus  $\mu(\{x_\beta : \beta > \alpha\}) = 0$ . Since  $\text{non}(\mathcal{N}) \geq \kappa$ ,

$$\mu(\{x_\beta : \beta \leq \alpha\}) = 0.$$

It follows that  $\mu(X) = 0$ .  $\square$

Given  $f: \omega \rightarrow \omega$  and superperfect  $T \subseteq \omega^{<\omega}$ , let  $f_T: \omega \rightarrow \omega$  be such that, whenever  $s$  is an  $n$ th-level splitnode of  $T$ , and  $s(k) \leq f(k)$  for all  $k < |s|$ ,  $f_T(n) > f(|s|)$ .

Then, whenever  $s$  is an  $n$ th-level splitnode of  $T$ , there is a  $k \leq |s|$  such that  $(s \frown \langle f_T(n) \rangle)(k) > f(k)$ .

Given  $F = \{f_\alpha : \alpha < \beta\}$  and  $T$ , let  $F'$  be the set of  $f \in \omega^\omega$  which agree with some  $f \in F$  mod-finite. Let  $h \in \omega^\omega$  be such that, for all  $f \in F'$ ,  $f_T \not\leq^* h$ .

Let  $x_\alpha$  extend infinitely many splitnodes of  $T$ , in such a way that, for each initial segment  $s$  of  $x_\alpha$ , if  $s$  is an  $n$ th level splitnode of  $T$ , then  $x_\alpha(n) = h(n)$ .

Does the existence of a nonmeager universally null set follow from either  $\mathfrak{d} = \text{cof}(\mathcal{M})$  or  $\text{non}(\mathcal{N}) \geq \text{cof}(\mathcal{M})$ ?

Note that if

$$\text{non}(\mathcal{N}) > \text{cof}(\mathcal{M})$$

then

$$\text{non}(\mathcal{N}) > \text{non}(\mathcal{M}),$$

in which case there are nonmeager universally null sets of cardinality  $\text{non}(\mathcal{M})$ . So the interesting case is when

$$\text{non}(\mathcal{N}) = \text{cof}(\mathcal{M}).$$

T. Bartoszyński, H. Judah, **Set theory**, A K Peters, 1995

D.P. Bertsekas, S.E. Shreve, **Stochastic Optimal Control: The Discrete-Time Case**, Academic Press, 1978

L. Bukovský, **The Structure of the Real Line**, Springer Science & Business Media, 2011

J.P. Burgess, *Classical hierarchies from a modern standpoint, Part I*, Fund. Math 115 (1983) 2, 81-95

J.P.R. Christensen, *Borel structures in groups and semigroups*, Math. Scand. 28 (1971), 124–128

A. S. Kechris, **Classical Descriptive Set Theory**, Springer-Verlag, 1995

A.S. Kechris, S. Solecki, S. Todorcevic, *Borel chromatic number*, Adv. Math. 141 (1999), no. 1, 1–44

R.D. Mauldin, D. Preiss, H. von Weizsäcker, *Orthogonal transition kernels*, Ann. Probab. 11 (1983), no. 4, 970–988

A. Miller, *Special subsets of the real line*, Handbook of Set-Theoretic Topology, North-Holland, 1984

I. Reclaw, *On a construction of universally null sets*, Real Anal. Exchange 27 (2001/02), no. 1, 321–323

C. Rosendal, *Continuity of universally measurable homomorphisms*, Forum Math. Pi 7 (2019)